

# Ethnomedicinal uses of plants among the Somali ethnic group, Jigjiga Woreda, Somali Regional State, Eastern Ethiopia

Arebu Issa Bilal,<sup>1</sup> Teferi Gedif Fenta,<sup>1</sup> Tsige Gebre-Mariam,<sup>1</sup> Kaleab Asres<sup>2\*</sup>

## Abstract

**Introduction:** In Ethiopia, most people are dependent on traditional medicine (TM), mainly of plant origin for human and animal health problems. The practice of herbal medicine varies widely, in keeping with the societal and cultural heritage of different countries. The heritage has not been well documented in the developing countries including Ethiopia and this is even more so in the emerging regions of the country. The objective of this study was to document medicinal plant knowledge of the people and identifying factors determining the use of medicinal plants in Jigjiga Woreda, Somali Regional State, eastern Ethiopia.

**Methods:** A cross-sectional study was conducted in five kebeles from where ethno-medicinal information was collected using semi-structured questionnaire. The questionnaire was administered to 800 heads of households. Data were entered and analyzed using Statistical Packages for Social Sciences version 20. Multivariable logistic regression was performed to show possible associations between the dependent and independent variable and statistical significance was set at  $p < 0.05$ .

**Results:** The study documented an overall prevalence of 40% to use of herbal medicine during one month recall period. A total of 45 medicinal plant species were collected and botanically identified. The study found that leaves are the most frequently utilized plant part (30.5%), followed by roots (23.0%). The reasons for preference of herbal drugs were related to lower price, efficacy and geographic accessibility as compared to modern medicine. Age, gender, educational status and occupation were identified as important determinants for the use of herbal medicine.

**Conclusion:** This ethno-medicinal study showed that community in Jigjiga Woreda relies on traditional medicinal plant species to treat a wide spectrum of human ailments. It is therefore suggested that more in depth studies be conducted to explore the potential of traditional medicine in the region to preserve this indigenous knowledge. [*Ethiop. J. Health Dev.* 2017;31(3):188-199]

**Key words:** Ethnomedicine, Jigjiga Woreda, Medicinal plants, Somali ethnic group

## Introduction

In Ethiopia, traditional mechanisms of dealing with illness have been there for centuries. The practice of traditional medicine (TM) to a large extent focuses on the use of herbs, spiritual healing, bone-setting, and minor surgical procedures. The practice of TM varies in its form, procedure, and content according to local customs (1). More than 95% of traditional medical preparations in Ethiopia are of plant origin (2). Plant-based TM plays a key role in the development and advancement of modern medicine by serving as a starting point for the development of novelties in drug discovery (3). In spite of increase in the health service coverage of the country, studies have reported that still traditional medicine plays important role in healthcare in Ethiopia (4, 5). Nevertheless, the loss of valuable medicinal plants due to lack of documentation, underreporting, agricultural expansion and deforestation, is widely reported by different researchers in Ethiopia (6-10). Available evidences suggest that studies on medicinal plants in Ethiopia have concentrated in the south, southwest, central, north and north-western parts of the country (4, 5, 11-36). There is little data available in the literature that assess the resource potential, indigenous knowledge

and preferences of use of medicinal plant species in eastern Ethiopia (37-39), and none of them are from the present study area.

Documentation of knowledge of medicinal plants is of great value to facilitate discovery of new sources of drugs and promote sustainable use of natural resources. Similarly, the knowledge of factors involved in the selection of treatment options at household (HH) level is important for health service planning and to incorporating herbal medicine in a country's health care delivery system (32). Thus, this study was carried out to document traditional medicinal plant knowledge of the people and factors determining the use of medicinal plants in Jigjiga Woreda, Somali Regional State, Eastern Ethiopia.

## Methods

**Study area:** Somali Regional State is one of the nine Federal States of Ethiopia located in the eastern part of the country with an estimated area of 279,252 km<sup>2</sup>. According to the 2007 population and housing census of Ethiopia, the total population of Somali Region was 4,445,219 which gives a population density of 15.9 /km<sup>2</sup> (40). Based on this census, Jigjiga Woreda

<sup>1</sup>Departement of Pharmaceutics and Social Pharmacy, School of Pharmacy, College of Health Sciences, Addis Ababa University, Ethiopia, E-mail Arebu – arebu.issa@aau.edu.et; Teferi - teferi.gedif@aau.edu.et; Tsige – tsige.gmariam@aau.edu.et; Kaleab - kaleab.asres@aau.edu.et

<sup>2</sup>Department of Pharmaceutical Chemistry and Pharmacognosy, School of Pharmacy, College of Health Sciences, Addis Ababa University, Ethiopia, E-mail kaleab.asres@aau.edu.et. P. O. Box 1176, Addis Ababa, Ethiopia

has a total population of 277,560; of those 125,876 (45.35%) are urban inhabitants, and 6,956 (2.51%) are pastoralists and the rest are engaged in crop farming, trade and non-farm related jobs. The woreda is primarily inhabited by the Geri tribe and small numbers of the Bartire (Yabaree) tribe. At the time of the survey, the Somali Region had 10 hospitals, 131 health centers and 950 health posts, of which 2 hospitals, 5 health centers and 26 health posts are located in Jigjiga Woreda (41).

**Sampling and data collection procedures:** Among the forty-six kebeles (the smallest administrative units) found in Jigjiga Woreda, 36 are rural and 10 are urban. After stratifying the kebeles into rural and urban, four rural kebeles and one urban kebele were selected based on probability proportional to population size. The kebeles selected were Ellebahay, Duhedi, Hadew, Qordehaere and Kebele 08. The number of household (HHs) heads included from each selected kebele was again determined based on the probability proportional to size and identified using systematic random sampling techniques where every fifth HH was taken until the required size was met in each kebele. The field work was carried out between January and March 2014. Data on medicinal plant use and preparation of plant specimens were collected. For each plant local Somali vernacular name, its uses or effects, the part(s) of the plant used, place of collection, its preparation and administration process were collected. At the end of the interviews, specimens of plants mentioned for medicinal uses were collected along with its specifications (vernacular names and the plant part used were recorded). Botanically identified plant specimens were stored at the National Herbarium, Addis Ababa University for further ethno pharmacological study.

**Sample size determination:** The minimum number of sample required for this study was determined by using single population proportion formula considering the following assumption

$$n_i = \frac{(Z\alpha/2)^2 p(1-p)}{d^2}$$

Where

$n_i$  = minimum sample size required for the study

Z = standard normal distribution (Z=1.96) with confidence interval of 95% and  $\alpha = 0.05$

P = the prevalence of TM use in Jigjiga Woreda, P = 50 % (0.5) will be used since there is no any prior study.

D = Absolute precision or tolerable margin of error (d) = 5% = 0.05

$$n_i = \frac{(Z\alpha/2)^2 p(1-p)}{d^2} = \frac{(1.96)^2 \times 0.5(1-0.5)}{(0.05)^2} = 384$$

To correct for the design effect “n” was multiplied by the number of stages, 2.  $N \times 2 = 768$

Considering 10% non-response rate, the final sample size was =845. Ten data collectors and one supervisor, who are health science students with knowledge of the local language, were employed and given training for five days on the data collection instruments.

**Inclusion criteria:** Household heads who were aged 18 or older, lived in the selected kebele for at least six months prior to the data collection, and gave informed consent were included in the study.

**Data collection instruments:** A pre-tested, semi-structured questionnaire was used for the data collection. The questionnaire was prepared in English and translated into the local language (Somali). The questionnaire contained information on demographic characteristics, history of illness in the family in the one month preceding the interview date, action(s) taken, reasons for the choice of herbal medicine as the first line of action and medicinal plants used.

**Ethical clearance:** The Institutional Ethical Review Board of the School of Pharmacy of Addis Ababa University cleared and approved to conduct the survey. Discussions about the aim and purpose of the survey were held with Jigjiga Woreda Health Office and local community leaders, who were asked for their cooperation. Oral consent was obtained from study participants for conducting interviews as well as taking their pictures for possible publication.

**Data entry and analysis:** Statistical Packages for Social Sciences (SPSS) version 20 was used for data entry and analysis of the data. Any logical and consistency errors identified during data entry were corrected after revision of the original completed questionnaire. Multivariable logistic regression was performed to show possible associations between the dependent (health-seeking behavior) and independent variables (age, gender, residence, household size of educational status and occupational status) and Statistical significance was set at  $p < 0.05$ .

## Results

**Use of medicinal plants:** A total of 45 medicinal plant species with reported medicinal values were collected and botanically identified during the course of this study (Table 1). The identified plants fall under 28 plant families with the largest number in the Solonaceae, followed by Euphorbiaceae and Asteraceae. According to the respondents, the most frequently utilized plant parts were leaves (30.5%), followed by root (23%), seeds (18%), and stems (7.5%). Plant parts such as fruits, rhizomes, flowers and bark are seldom used (Table 1).



Table 1: **Plants used in the treatment of human disorders in Jiggiga Woreda, March 2014**

Scientific name (Collection number)	Family	Vernacular name	Sign and symptoms (number of responses)	Part used	Method of preparation and administration
<i>Acokanthera schimperi</i> (A.DC.) Schweinf (AI-41)	Apocynaceae	Qarari	<i>Shuban</i> (1)	Seed	Ground, dispersed in water and drunk
<i>Allium sativum</i> L. (AI-15)	Alliaceae	2. Toon	<i>Qufac</i> (13), <i>Duray</i> (5) <i>Il shaydaan</i> (2) <i>Wadna xanuun</i> (1) <i>Feex</i> (1) <i>Laab wareen</i> (1) <i>Xumad</i> (1) <i>Ilka xanuun</i> (1) <i>Xanuun kaste</i> (1)	Bulb	Crushed, mixed with tea and drunk Ground, mixed with milk and sugar, boiled and drunk Chewed
<b><i>Aloe megalacantha Baker</i></b> (AI-33)	Aloaceae	Dacar	<i>Indha xanuun</i> (17) <i>Wadna xanuun</i> (2) <i>Calool ingeeg</i> (3) <i>Infekshinka hunguriga</i> (2) <i>Indha xanuun</i> (1) <i>Laqanyo</i> (1) <i>Matag</i> (1)	Leaf Latex	Pulverized in water and applied on the affected area Dissolved in water and drunk
<i>Alternanthera pungens</i> Kunth. (AI-5)	Amaranthaceae	Gucundho	<i>Barar</i> (3) <i>Kadiolol</i> (2)	Whole part	Crushed, boiled with water and applied on the affected area
<i>Amaranthus caudatus</i> L. (AI-22)	Amaranthaceae	Milaxbuur	<i>Xanuun kaneefita</i> (2) <i>Qixdheer</i> (1) <i>Barar</i> (1)	Leaf	Pounded and eaten with food Crushed in boiled water sugar added, and drunk
<i>Azadirachta indica</i> A. Juss. (AI-8)	Meliaceae	Geed hindi	<i>Dhabar xanuun</i> (1) <i>Ilka xanuun</i> (5) <i>Calool guruuruc</i> (2)	Leaf	Boiled and decoction drunk
<i>Brassica oleracea</i> L. (AI-13)	Brassicaceae	Sagaxoor	<i>Xanuun kaneefita</i> (2) <i>Qix dheer</i> (1) <i>Barar</i> (1)	Leaf	Crushed and eaten with food or boiled with sugar and drunk
<i>Capsicum annum</i> L. (AI-9)	Solanaceae	Filfil	<i>Kabiibeyso</i> (1)	Seed	Chewed and placed on tooth
<i>Carica papaya</i> L. (AI-14)	Caricaceae	Papyee	<i>Dirxiga mindhiqirka</i> (1)	Seed	Pounded, dispersed in water and drunk
<i>Casimiroa edulis</i> La Llave (AI-43)	Rutaceae	Canbe	<i>Cagaarshow/joonis</i> (1)	Leaf	Boiled and decoction drunk with sugar

<i>Citrus limon</i> (L.) Burm.f. AI-11	Rutaceae	Lendenan	<i>Shuban</i> (1)	Fruit	Squeezed and the juice drunk
<i>Coffea arabica</i> L. (AI-40)	Rubiaceae	Quid	<i>Shuban</i> (1) <i>Dhiig laaan</i> (1)	Seed	Pounded, boiled for three hours and drunk
<i>Combretum molle</i> R. Br. Ex. G. Don (AI-24)	Combretaceae	Woob	<i>Faalid</i> (1) <i>Beer xanuun</i> (5) <i>Barar</i> (1)	Root Bark	Crushed, mixed with oil and applied to the affected area Pounded, dispersed in water and drunk
<i>Coriandrum sativum</i> L. (AI-6)	Apiaceae	Qorjeen	<i>Dhabar xanuun</i> (15) <i>Roomatism</i> (8)	Latex Seed	Boiled and decoction drunk Crushed, mixed with honey or sugar and drunk or eaten with food
<i>Croton macrostachyus</i> Del. (AI-37)	Euphorbiaceae	Mekenisa	<i>Beer xanuun</i> (2)	Bark	Pounded, mixed with water and drunk
<i>Euphorbia polycantha</i> Boiss. (AI-18)	Euphorbiaceae	Waantays	<i>Gaaska</i> (2)	Leaf	Crushed, dispersed in water and drunk
<i>Euphorbia tirucalli</i> L. (AI-12)	Euphorbiaceae	Iin	<i>Feex</i> (1)	Latex	Dissolved in water, sugar added and drunk
<i>Jasminum grandiflorum</i> L. subsp. <i>floribundum</i> (R.Br.ex Fresen.) P.S.Green (AI-35)	Oleaceae	Qajajuli	<i>Boog</i> (1)	Root	Pounded and placed on wound
<i>Lepidium sativum</i> L. (AI-20)	Brassicaceae	Shunfax	<i>Qufac</i> (11) <i>Dhabar xanuun</i> (7) <i>Laab wareen</i> (9) <i>Cudurka qaaxada</i> (6) <i>Ilka xanuun</i> (2) <i>Shuban</i> (2) <i>Indha xanuun</i> (1) <i>Xanuun kaste</i> (1)	Seed	Fresh seeds swallowed
<i>Lippia adoensis</i> Hochst. ex Walp. (AI-42)	Verbenaceae	Sokay	<i>Jiljilee ccaruurta iyo hooyada uurkaleh</i> (1) <i>Sanboor</i> (4)	Leaf	Crushed, boiled in water with sugar and drunk
<i>Melia azedarach</i> L. (AI-19)	Meliaceae	Geed kinin	<i>Ilka xanuun</i> (1)	Leaf	Pounded and pressed on tooth

<i>Moringa stenopetala</i> (Bak. F.) Cuf. (AI-3)	Moringaceae	Moringa	<i>Xanuunka Macaan (1)</i> <i>Dhiig kar (1)</i> <i>Dhiig laaan (1)</i> <i>Loo isticmaalo qurxinta</i> <i>Jidhka (1)</i>	Leaf	Ground, dispersed in tea and drunk
<i>Nicotiana glauca</i> R.Grah. (AI-45)	Solanaceae	Booc	<i>Infekshinka maqaarka (1)</i>	Leaf	Squeezed between palms, the fluid mixed with oil and applied on the affected area
<i>Ocimum lamiifolium</i> Hochst. ex Benth. (AI-34)	Lamiaceae	Demakesae	<i>Qufac (1)</i>	Leaf	Crushed, squeezed, and a teaspoon full drunk with tea or coffee
<i>Olea europaea</i> L. subsp. <i>cuspidate</i> (WalL.exG.Don) Cif (AI-16)	Oleaceae	Ejersa	<i>Finanka (1)</i> <i>Barse (1)</i>	Root	Pounded, mixed with oil and applied on the affected area
<i>Osyris quadripartite</i> Decn. (AI-25)	Santalaceae	Wato	<i>Duumad (Kaneeco) (1)</i>	Leaf	Macerated for 24 h and the reddish macerate drunk
<i>Parthenium hysterophorus</i> L. (AI-1)	Asteraceae	Kalignoole	<i>Dhiig bax (18)</i> <i>Teetano (1)</i>	Leaf	Pounded and placed on wound
			<i>Dhabar xanuun (1)</i> <i>Ilka xanuun (1)</i>	Root	Chewed and pressed on tooth
<i>Prunus persica</i> (L.) Batsch (AI-38)	Rosaceae	Kuki	<i>Beer xanuun (1)</i>	Leaf	Steeped in cold water and infusion drunk
<i>Psidium guajava</i> L. (AI-36)	Myrtaceae	Saytuun	<i>Shuban (1)</i> <i>Matag (1)</i>	Leaf	Crushed in water and drunk with sugar
<i>Punica granatum</i> L. (AI-27)	Lythraceae	Ruman	<i>Il shaydaan (2)</i>	Leaf/ Seed	Crushed leaves and seeds mixed with milk and drunk
<i>Rhamnus prinoides</i> L'Herit. (AI-44)	Rhamnaceae	Gesho	<i>Calool ingeeg (1)</i>	Leaf	Pulverized in cold water and drunk
<i>Ricinus communis</i> L. (AI-4)	Euphorbiaceae	Qobo	<i>Calool ingeeg (1)</i>	Root	Pounded, added to tea and drunk
<i>Ruta chalepensis</i> L. (AI-23)	Rutaceae	Taltan	<i>Il shaydaan (3)</i>	Leaf	Boiled and decoction drunk

<i>Schinus molle</i> L. (AI-26)	Anacardiaceae	Mirmir	<i>Dirxi</i> (3) <i>Gooriyaan</i> (1) <i>Dirxi noocbalaaran</i> (1) <i>Ilka xanuun</i> (1) <i>Calool ingeeg</i> (1)	Leaf	Crushed, dispersed in water and drunk or powder pressed on tooth
<i>Senna italica</i> Mill. (AI-21)	Fabaceae	Jaleelo		Leaf	Pounded, mixed with water and drunk
<i>Solanecio angulatus</i> (Vahl) C, Jeffrey (AI-32)	Asteraceae	Hadus	<i>Duaray</i> (1) <i>Xubno xanuun</i> (1) <i>Xumad</i> (1)	Leaf	Crushed, mixed with water and sugar and drunk
<i>Solanum giganteum</i> Jacq. (AI-29)	Solanaceae	Kariir	<i>Dhiigbax</i> (1)	Leaf	Crushed and applied on the wound
<i>Solanum incanum</i> L. (AI-28)	Solanaceae	Kariir	<i>Dhiigbax</i> (2)	Leaf	Pounded and applied on wound
<i>Solanum jubae</i> Bitter (AI-30)	Solanaceae	Kariir	<i>Dhiigbax</i> (1)	Leaf Root Bark	Mixed, ground and applied on wound
<i>Sphaeranthus suaveolens</i> (Forssk.) DC. (AI-31)	Asteraceae	Rashaid	<i>Maskax xanuun</i> (1)	Leaf Flower	Ground, mixed with water, oil and honey, and applied to the head
<i>Vernonia amygdalina</i> Del. (AI-39)	Asteraceae	Girawae	<i>Dirxiga mindhiqirka</i> (1) <i>Barar</i> (1) <i>Il shaydaan</i> (1)	Leaf	Ground, dispersed in water and drunk
<i>Viscum tuberculatum</i> A. Rich. (AI-7)	Viscaceae	Dhigri	<i>Shuban</i> (10) <i>Laqanyo</i> (8) <i>Xumad</i> (4) <i>Madax xanuun</i> (2) <i>Dhiigbax</i> (1)	Root	Pounded, boiled and decoction drunk
<i>Withania somnifera</i> (L.) Dunal (AI-17)	Solanaceae	Kariir		Leaf	Powdered and placed on wound
<i>Zingiber officinale</i> Roscoe (AI-10)	Zingiberaceae	Singibill	<i>Qufac</i> (4) <i>Duaray</i> (4) <i>Quman/qanjiroxanuun</i> (1) <i>Infekshinka hunguriga</i> (1) <i>Nadaafada wajiga iyo daawaynta finanka wajiga</i> (1)	Bulb	Crushed and decoction drunk with sugar or powder drunk with tea
<i>Ziziphus mauritiana</i> Lam. (AI-2)	Rhamnaceae	Kasil		Leaf	Pounded, stood in cold water and applied on the face

**Collection and storage of plant medicines:** The major proportion of plants collected by HHs is from wild sources (70%); only 20% is cultivated and the remaining 10% is from both sources. Most of the respondents stored herbal formulations in plastic bags, tin containers, bottles and paper bags. Route of administration.

The most widely used route of administration was found to be oral (66.8%), followed by topical application (19.5%), oral and topical (3.6%), and as eye drops (2.2%). Lack of precision with doses has been noted where 70% reported to approximate while (19.8%), use cups, (6%) tea spoons and (3%) hands.

**Perceived illness and health-seeking behavior:** The proportion of respondents with perceived illness during one month recall period preceding the interview date was 251(31.3%). Females 127 (50.5%) had more

morbidity than males 124 (49.4%) and those aged between 15- 65 had more frequent morbidity 101 (40.2%), but the age group of greater than 65 had less morbidity 5 (1.9%) than the other age groups.

In response to perceived illness episodes, a large proportion 241 (96.0%) of those who reported illness took action against their illness. While 134 (53.3%) of them went to health institutions, 85 (33.8%) visited local healers, 14 (5.5%) used homemade herbal remedies, while 10 (4.0%) took no action (Table 2). Therefore, 40% of the respondents in this study reported to use traditional medicine during the recall period. Most (62.7%) of the urban residents visited health facility while 28.9% of them consulted local traditional healers. In rural setting however large proportion (45.1%) of patient consulted local traditional healers while 36.5% of them visited health facility (Table 2).

**Table 2: Types of action taken by those with perceived illness during the one month recall period (N = 251), in Jigjiga Woreda, March, 2014**

Factor	Consulted healers N (%)	Went to health facility N (%)	Used homemade remedies N (%)	No action taken N (%)	Both healers And health Facilities N (%)	Total
<b>Gender</b>						
Male	48 (38.7)	62 (50)	7 (5.6)	4 (3.2)	3 (2.4)	124 (49.4)
Female	38 (30)	72 (56.6)	7 (5.5)	6 (4.7)	4 (3.0)	127 (50.5)
<b>Age</b>						
<5	18 (23.0)	48 (61.5)	6 (7.6)	5(3.8)	1(1.2)	78 (31.0)
5-15	27 (40.2)	32 (47.7)	5 (7.4)	2 (2.9)	1(1.4)	67 (26.6)
15-65	41(40.5)	53 (52.4)	4 (3.9)	3 (2.9)	0	101 (40.2)
>65	2 (40)	2 (40)	1(20)	0	0	5 (1.9)
<b>Residence</b>						
Urban	49 (28.9)	106 (62.7)	6 (3.5)	5 (3.5)	3 (1.7)	169 (67.4)
Rural	37 (45.1)	30 (36.5)	8 (9.7)	5 (6.0)	4 (4.8)	82 (32.6)
<b>Total</b>	86 (34.2)	133 (53)	14 (5.5)	10 (4.0)	8 (3.0)	251 (100)

Result from multivariate logistic regression shows that males were found to use modern health institution (AOR=1.55; 95%CI [0.89-2.70]) as compared to women although the association was not statistically significant. As age increases however, the probability of using modern health institutions decreases (AOR=0.73; 95% CI [0.35-1.54]); (AOR=0.23; 95%CI

[0.07-0.71]). Residence showed a statistically significant association with the use of modern health institutions, where urban residents were found to use modern health institution more than twice as compared to rural residents (AOR=2.80; 95%CI [1.54-5.09]) (Table 3).

Table 3: Factors affecting patterns of health-seeking behavior among household respondents in Jigjiga Woreda, during the one month recall period [March 2014 (N = 251, 95% Confidence interval)].

	Traditional medicine	Modern health institution	Crude odds ratio	Adjusted odds ratio
<b>Gender</b>				
Male	55	62	1.00	1.00
Female	45	72	1.14 [0.84-2.38]	1.55 [0.89-2.70]
<b>Age</b>				
6 months-25	66	110	1.00	1.00
25-45	20	19	0.60 [0.29-1.21]	0.73 [0.35-1.54]
Greater than 65	14	5	0.21 [0.07-0.62]*	0.23 [0.07-0.71]*
<b>Residence</b>				
Rural	45	29	1.00	1.00
Urban	55	105	2.89 [1.64-5.08]*	2.80 [1.54-5.09]*

\* Statistically significant

**Factors associated with health-seeking behavior:**

Preference of healthcare was assessed by asking about usual health-seeking practices during illness. As shown in Table 4, 684 (85.5%) of the respondents preferred to go to health institutions, 99 (12.3%) preferred healers, and 17 (2.2%) preferred to use homemade herbal remedies. More females (13.6%) preferred visiting THs than males (10.0%).

Reasons for preference of TM shows that low price 75 (65.5%), effectiveness 25 (21.5%) and access 15 (13.0%) were the major reasons for preference of traditional medicine. Age groups greater than 65 years

preferred to use TM as compared to respondents with the age group 18-25 years (AOR = 3.473; 95%CI [1.188-10.155]). The influence of education on treatment preference was analyzed and it was found that non literates tended to use TM more than literates and the relationship was statistically significant, (AOR = 2.337; 95% CI [1.438-3.798]). Residence of respondents did not show a significant association with the preferences of TM use, (AOR=0.85; 95% CI [0.48-1.49]). Use of TM was viewed negatively more among housewives and merchants than agro-pastoralists (AOR =0.42; 95%CI [0.222-0.789]) and (AOR = 0.220; 95%CI [0.06-0.72] (Table 3).

Table 4: Factors affecting patterns of health-seeking behavior among household respondents in Jigjiga Woreda, March 2014 (N = 800, 95% Confidence interval).

	Traditional medicine	Modern health institution	Crude odds ratio	Adjusted odds ratio
<b>Gender</b>				
Female	80	431	1.30 [0.85-1.99]	*1.76 [1.07-2.92]
Male	36	253	1.00	1.00
<b>Residence</b>				
Rural	58	281	1.43 [0.967-2.128]	0.85 [0.48-1.49]
Urban	58	403	1.00	1.00
<b>Age</b>				
18-25	8	89	1.00	1.00
26-45	75	414	2.01 [0.93-4.32]	1.83 [0.82-4.05]
46-65	23	149	1.71 [0.73-4.00]	1.579 [0.65-3.82]
>65	10	32	*3.47 [1.26-9.58]	*3.47 [1.18-10.15]
<b>Household size</b>				
1-4	19	133	1.00	1.00
5-8	73	394	1.29 [0.75-2.22]	1.090 [0.61-1.93]
>8	24	157	1.07 [0.56-2.03]	0.876 [0.44-1.73]
<b>Educational status</b>				
Literate	24	272	1.00	1.00
Non-literate	92	410	*2.54 [1.58-4.08]	*2.33 [1.43-3.79]
<b>Occupation</b>				
Agro-pastoralist	45	177	1.00	1.00
Housewife	47	311	*0.59 [0.38-0.93]	*0.41 [0.22-0.78]
Jobless	8	53	0.59 [0.26-1.33]	0.45 [0.19-1.08]
Merchant	4	63	*0.25 [0.08-0.72]	*0.22 [0.06-0.72]
Government employee	1	19	0.20 [0.02-1.58]	0.24 [0.02-2.04]
Others*	11	61	0.70 [0.34-1.45]	0.70 [0.28-1.73]
<b>Total</b>	116 (14.5%)	684 (85.5%)		

\* Statistically significant; \*\* Butchers, wood collectors, shoe makers

## Discussion

The present study revealed that the overall proportion of reported illness episodes among the Somali ethnic group in Jigjiga Woreda during the one month recall period was 31.3%. Although different studies documented variations in prevalence of illness episodes during the recall periods, most study documented widespread use of TM under illness circumstances (4, 33, 34). A study conducted in Addis Ababa found 37% of study participants use herbal medicine (33), and studies carried out in other parts of Ethiopia such as Wombera District of Benishangul-Gumuz Region reported 41.6% (4) of respondents preferred traditional medicine. In the current study, 40.0% of participants reported to use traditional medicine. Studies conducted in Butajira and among the Berta ethnic group found 12.5% and 4.6% use of traditional medicine in four weeks and two weeks recall period, respectively (5, 34). This might be due to differences in the literacy status and health service coverage between the regions (40, 41). Reasons given for preferring herbal drugs were lower price, better efficacy and inaccessibility to modern medicine. The study also showed that more females (13.0%) prefer visiting THs than males (10.0%). Similar findings were reported in earlier works carried out in other parts of Ethiopia (5, 34, 35).

In this study, the preference of TM by housewives and merchants was negatively associated, (AOR = 0.42; 95% CI [0.22- 0.79]), AOR= 0.22; 95% CI [0.06-0.72]), respectively) as compared to the use by agro-pastoralists. As agro-pastoralists spend most of their time in the field, they may have better knowledge about TM and herbs than people in other sectors (42). The study also showed that people older than 65 years and illiterate preferred TM significantly more than younger age groups. A similar finding was reported in different parts of the country where illiterates and older residents were significantly more likely to use herbal medicine than literate and younger people (33, 34, 36, 37, 39). Other studies indicated that younger generation are less knowledgeable about TM and tend to underestimate traditional values (39, 43, 44). In contrast to the above findings, a study conducted in Suriname found that age and educational status were not the predictors of herbal medicine use (45).

According to the findings of the present study, leaves are the most frequently utilized plant part (30.5%), followed by roots (23.0%) which is similar to previous studies conducted in other parts of the country (6, 37, 39, 47,48). However, in some other studies, roots have

been reported to be the most extensively used part of medicinal plant (4, 8, 26, 34, 49, 50). It is well known that herbal preparations that involve roots, rhizomes, bulbs, barks, stems or whole parts have effects on the survival of the mother plant (51). The possible destruction of medicinal plants due to collection for medicinal purposes is minimal since leaves were the leading plant part used in the area. However, since most of the medicinal plants reported are obtained from the wild (70.0%) this can pose threat to biodiversity. The study showed that there is lack of precision in the determination of doses of medicaments used, which is consistent with other works conducted in different parts of the country (34, 37, 39, 50). The real drawback in TM mostly arises from lack of precision in dosage (51).

#### **Conclusion:**

This ethnomedicinal study showed that rural and urban communities in Jigjiga Woreda still rely on medicinal plants to treat a wide spectrum of human ailments. A total of 45 medicinal plant species along with their uses have been reported, and the majority of them are harvested from the wild. The study found that leaves are the most frequently utilized plant part, followed by roots. The reasons for preferring herbal drugs were lower price, efficacy and geographic inaccessibility to modern medicine. Age, gender, educational status and occupation are important factors that determine the use of TM in the study area. Many young people, males, literate and occupations other than agro-pastoralists were found to have less preference for the use of TM for their illness. It is, therefore, suggested that other studies need to be conducted to explore the potential of the different Woredas and zones of the region to preserve this indigenous knowledge of TMs by proper documentation, identification of plant species, herbal preparation and dosage forms used.

#### **Acknowledgements**

Our appreciation goes to the people of Jigjiga Woreda, and community leaders for their positive response, sharing their valuable knowledge and time as well as for their tremendous generosity and hospitality. The authors also would like to acknowledge School of Pharmacy, College of Health Sciences, Addis Ababa University (AAU), for sponsoring this research work which is part of an MSc study. The authors would also like to thank Mr. Muktar Sheh Abdulahi, Bedrie Abdulahi, Ibrahim Dawud, Hirute, Hikmat Mohammed and Abduwahab for their tireless support in the translation and collection of specimen. We would also like to extend our gratitude to Prof. Sebsebe Demissew, Mr. Melaku Wondafrash and Mr. Wegae Abebe, the National Herbarium, Department of Biology, AAU, for identification of plant specimens.

#### **Author Disclosure Statement**

The authors declare that they have no competing financial interests.

#### **Authors' contributions**

AI coordinated data collection; performed data entry and analysis; wrote the draft manuscript. TG, KA and TGM initiated the idea; involved in the design of the study; developed data collection instruments and corrected the manuscript. All the authors have read and approved the final manuscript.

#### **References**

1. Pankhurst R. An introduction to the medical history of Ethiopia. Trenton, New Jersey: The Red Sea Press, Inc; 1990:68.
2. Abebe D. Traditional medicine in Ethiopia: the attempts being made to promote it for effective and better utilization. *SINET* 1986;9:61–69.
3. Wright CW: Plant derived antimalarial agents: new leads and challenges. *Phytochemistry* 2005;4:55–61.
4. Guji T, Gedif T, Asres K, Gebre-Mariam T. Ethnopharmaceutical study of medicinal plants of Metekel Zone Benishangul–Gumuz regional state mid-west Ethiopia. *Ethiop Pharm J* 2011;29:43–58.
5. Gedif T, Hahn H. The use of medicinal plants in self-care in rural central Ethiopia. *J Ethnopharmacol* 2003;87:155–161.
6. Abebe D. Biodiversity conservation of medicinal plants: Problem and prospects. In *Conservation and sustainable use of medicinal plants in Ethiopia. Proceeding of The National Workshop on Biodiversity Conservation and Sustainable Use of Medicinal Plants in Ethiopia* Edited by: Zewdu M, Demissie A. Addis Ababa: IBCR; 2001:198–203.
7. Getachew B, Shiferaw D. Medicinal plants in Bonga Forest and their uses. *Biodiversity Newslett* 2002;1:9–10.
8. Addis G, Abebe D, Urga K. A survey of traditional medicinal plants in Shirka district, Arsi zone, Ethiopia. *Ethiop Pharm J* 2001;19:30–47.
9. Yineger H, Yewhalaw D. Traditional medicinal plant knowledge and use by local healers in Sekoru District, Jimma Zone, Southwestern Ethiopia. *J Ethnobiol Ethnomed* 2007;3:24.
10. Birhanu Z. Traditional use of medicinal plants by the ethnic groups of Gondar Zuria District, north-western Ethiopia: *J Nat Rem* 2013;13:2320–3358.
11. Giday M, Teklehaymanot T. Ethnobotanical study of plants used in management of livestock health problems by Afar people of Ada'ar district, Afar regional state, Ethiopia. *J Ethnobiol Ethnomed* 2013;9:8.
12. Giday M, Asfaw Z, Thomas E, Woldu Z. An ethnobotanical study of medicinal plants used by the Zay people in Ethiopia. *J Ethnopharmacol* 2003;85:43–52.
13. Tolossa K, Debela E, Athanasiadou S, Tolera A, Ganga G, Jos GMH. Ethno-medicinal study of plants used for treatment of human and livestock ailments by traditional healers in South Omo, Southern Ethiopia. *J Ethnobiol Ethnomed* 2013;9:32.
14. Birhan W, Giday M, Teklehaymanot T. The contribution of traditional healers' clinics to public

- health care system in Addis Ababa, Ethiopia: a cross-sectional study. *J Ethnobiol Ethnomed* 2011;7:39.
15. Yineger H, Yewhalaw D. Traditional medicinal plant knowledge and use by local healers in Sekoru District, Jimma Zone, Southwestern Ethiopia. *J Ethnobiol Ethnomed* 2007;3:24.
  16. Giday M, Teklehaymanot T, Animut A, Mekonnen Y. Medicinal plants of the Shinasha, Agew-Awi and Amhara peoples in northwest Ethiopia. *J Ethnopharmacol* 2007;110:516–525.
  17. Giday M, Asfaw Z, Woldu Z. Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *J Ethnopharmacol* 2010;132(1):75–85.
  18. Abbink J. Me'en ritual, medicinal and other plants: a contribution to southwest Ethiopian ethnobotany. *J Ethnopharmacol* 1993;24:1–20.
  19. Balemie K, Kelbessa E, Asfaw Z. Indigenous medicinal utilization, management and threats in Fentale area, Eastern Shewa, Ethiopia. *Ethiop J Biol Sci* 2004;3:1–7.
  20. Yineger H, Yewhalaw D, Teketay D. Ethnomedicinal plant knowledge and practice of the Oromo ethnic group in southwestern Ethiopia. *J Ethnobiol Ethnomed* 2008;4:11.
  21. Bekalo T, Demissew S, Asfaw Z. An ethnobotanical study of medicinal plants used by local people in the lowlands of Konta special woreda, Southern Nations, Nationalities and Peoples Regional State, Ethiopia. *J Ethnobiol Ethnomed* 2009;5:26.
  22. Lulekal E, Kelbessa E, Bekele T, Yineger H. An ethnobotanical study of medicinal plants in Mana Angetu district, south-eastern Ethiopia. *J Ethnobiol Ethnomed* 2008;4:10.
  23. Yirga G. Use of traditional medicinal plants by indigenous people in Mekele town, capital city of Tigray regional state of Ethiopia. *J Med Plants Res* 2010;4:1799–1804.
  24. Zerabruk S, Yirga G. Traditional knowledge of medicinal plants in Gindeberet district, Western Ethiopia. *S Afr J Bot* 2012;78:165–169.
  25. Kloos H, Tekle A, Yohannes L, Yosef A. Preliminary studies of traditional medicinal plants in nineteen markets in Ethiopia: use patterns and public health aspects. *Ethiop Med J* 1978;16:33–43.
  26. Mesfin F, Demissew S, Teklehaymanot T. An Ethnobotanical study of medicinal plants in Wenago Woreda, SNNPR, Ethiopia. *J Ethnobiol Ethnomed* 2009;5:28.
  27. Teklehaymanot T, Giday M. Ethnobotanical study of medicinal plants used by people in Zegje peninsula, northwestern Ethiopia. *J Ethnobiol Ethnomed* 2007;3:12.
  28. Lulekal E, Asfaw Z, Kelbessa E, Van Damme P. Ethnomedicinal study of plants used for human ailments in Ankober District, North Shewa Zone, Amhara region, Ethiopia. *J Ethnobiol Ethnomed* 2013;9:63.
  29. Megersa M, Asfaw Z, Kelbessa E, Beyene A, Woldeab B. An ethnobotanical study of medicinal plants in Wayu Tuka district, east Welega zone of Oromia Regional State, West Ethiopia. *J Ethnobiol Ethnomed* 2013;9:68.
  30. Bussmann RW, Swartzinsky P, Worede A, Evangelista P. Plant use in Odo-Bulu and Demaro, Bale region, Ethiopia. *J Ethnobiol Ethnomed* 2011;7:28.
  31. Wondimu T, Asfaw Z, Kelbessa E. Ethnobotanical study of medicinal plants around “Dheeraa” town, Arsi zone Ethiopia. *J Ethnopharmacol* 2007;112:152–161.
  32. Birhane E, Aynekulu E, Mekuria W, Endale D. Management, use and ecology of medicinal plants in the degraded dry lands of Tigray, northern Ethiopia. *J Med Plants Res* 2011;5:309–318.
  33. Gedif T, Hahn HJ. Epidemiology of herbal drugs use in Addis Ababa, Ethiopia. *Pharmacoepidemiol Drug Saf* 2002;11:587–591.
  34. Flatie T, Gedif T, Asres K, Gebre-Mariam T. Ethnomedicinal survey of Berta ethnic group Assosa zone, Benishangul-Gumuz Regional State, mid-west Ethiopia. *J Ethnobiol Ethnomed*. 2009;5:14.
  35. Wolde B, Gebre-Mariam. Household herbal remedies for self-care in Addis Ababa: A preliminary assessment. *Ethiop Pharm J* 2002;20:59–70.
  36. Kidane B, Andel T, Maesen J and Asfaw Z. Use and management of traditional medicinal plants by Maale and Ari ethnic communities in southern Ethiopia. *J Ethnobiol Ethnomed* 2014;10:46.
  37. Belayneh A, Asfaw Z, Demissew S, Bussa N. Potential and use of medicinal plants by pastoral and agro-pastoral communities in Erer valley of Babile wereda, Eastern Ethiopia. *J Ethnobiol Ethnomed* 2012;8:42.
  38. Mesfin A, Giday M, Animut A, Teklehaymanot T. Ethnobotanical study of antimalarial plants in Shinile district, Somali region, Ethiopia, and *in vivo* evaluation of selected ones against *Plasmodium berghei*. *J Ethnopharmacol* 2012;139:1.
  39. Belayneh A, Bussa N. Ethnomedicinal plants used to treat human ailments in the prehistoric place of Harla and Dengego valleys, eastern Ethiopia. *J Ethnobiol Ethnomed* 2014;10:18.
  40. Ethiopia Central Statistical Authority Population and Housing Census Report-Somali Region - 2007. <http://www.csa.gov.et/index.php/2013-02-20-14-51-51/2013-04-01-11-53-00/census-2007>. Accessed on 15 March 2015.
  41. Federal Democratic Republic of Ethiopia Ministry of Health: Health and Health Related Indicators. Addis Ababa, Ethiopia; 2012. <http://www.moh.gov.et/documents/26765/0/Health+and+Health+Related+Indicators+2005+E.C/1b5b2a9f-a960-4024-8d92-519195364023?version=1.0>. Accessed on 10 January 2016.
  42. Feyera T, Mekonnen E, Wakayo BU, Assefa S. Botanical ethnoveterinary therapies used by agro-pastoralists of Fafan zone, Eastern Ethiopia. *BMC veterinary research*. 2017 Aug 9;13(1):232.
  43. Pankhurst R. The status and availability of oral and written knowledge on traditional health care in Ethiopia. In *Proceedings of the National*

- Workshop on Biodiversity Conservation and Sustainable Use of Medicinal Plants in Ethiopia Addis Ababa: IBCR; 2001:92–106.
44. Tanto T, Giday M, Aklilu N. National Biodiversity Strategy and Action Plan (BSAP) Project: Resource Base of Medicinal Plants of Ethiopia. Institute of Biodiversity Conservation 2002, Addis Ababa.
45. Van Andel T, Carvalheiro L. Why urban citizens in developing countries use traditional medicines? The case of Suriname. *Evid Based Complement Altern Med* 2013; 687197 doi: 10.1155/2013/687197.
46. Seifu T, Asres K, Gebre-Mariam T. Ethnobotanical and ethnopharmaceutical studies on medicinal plants of Chifera district, Afar region north eastern Ethiopia. *Ethiop Pharm J* 2006;24: 41–58.
47. Tadesse M, Hunde D, Getachew Y. Survey of medicinal plants used to treat human diseases in Sheka Chekorsa, Jimma zone, Ethiopia. *Ethiop J Health Sci* 2005;15:89–107.
48. Hunde D, Asfaw Z, Kelbessa E. Use of traditional medicinal plants by people of Boosat sub district, central eastern Ethiopia. *Ethiop. J Health Sci* 2006;16:141–155.
49. Giday M, Asfaw Z, Woldu Z. Medicinal plants of the Meinit ethnic group of Ethiopia: An ethnobotanical study. *J Ethnopharmacol* 2009;124;513–21.
50. ETANA T. Use and conservation of traditional medicinal plants by indigenous people in Gimbi Woreda, Western Wellega, Ethiopia. Accessed from T ETANA - 2007 - etd.aau.edu.et.
51. Abebe D, Ayehu A. Medicinal plants and enigmatic health practice of north Ethiopia. Berhanina Selam Printing Enterprise, Addis Ababa. 1993:34.